

WOODVALE PARK

VETERINARY HOSPITAL



Unit 10, 923 Whitfords Avenue, Woodvale WA 6026

Phone: (08) 9409 6968

www.woodvaleparkvet.com.au

Aimeroy Pty Ltd ABN 53 165 893 701

Pulmonic Stenosis

(Type of Heart Birth Defect)

Basics

OVERVIEW

- The heart of the dog or cat is composed of four chambers; the top two chambers are the right and left atria and the bottom two chambers are the right and left ventricles; heart valves are located between the right atrium and the right ventricle (tricuspid valve); between the left atrium and the left ventricle (mitral valve); from the right ventricle to the main pulmonary (lung) artery (pulmonary valve); and from the left ventricle to the aorta (the main artery of the body; valve is the aortic valve)
- “Pulmonic” refers to pulmonary; “pulmonary” refers to the lung(s); “pulmonic stenosis” is a condition characterized by narrowing of the pulmonary valve in the heart, which is the valve from the right ventricle to the main pulmonary artery
- “Stenosis” is the medical term for narrowing
- “Pulmonic stenosis” is a congenital (present at birth) narrowing at some point in the area through which blood flows out of the right ventricle, through the pulmonary valve, and into the main pulmonary artery (main artery of the lungs); this area is known as the “right ventricular outflow tract”; the narrowing blocks or obstructs the flow of blood from the right ventricle to the pulmonary artery
- Defect can be at the valve itself (known as “valvular pulmonic stenosis”), below the valve (known as “subvalvular pulmonic stenosis,” or above the valve, just inside the pulmonary artery (known as “supravalvular pulmonic stenosis”); valvular pulmonic stenosis is the most common form

GENETICS

- Inherited defect in beagles; multiple genes probably are involved (known as a “polygenic mode of transmission”)

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs
- Cats

Breed Predispositions

- English bulldog, Scottish terrier, wirehaired fox terrier, miniature schnauzer, West Highland white terrier, Chihuahua, Samoyed, mastiff, cocker spaniel, beagle, boxer

Mean Age and Range

- Present from birth and may be detected as a heart murmur in puppies
- If a heart murmur is not detected, affected pets may not be identified until clinical signs develop later in life

Predominant Sex

- Males are more likely than females to have pulmonic stenosis in English bulldogs and possibly other breeds

SIGNS/OBSERVED CHANGES IN THE PET

- Mild narrowing (stenosis)—usually no clinical signs
- Severely affected pets—may develop congestive heart failure (condition in which the heart cannot pump an adequate volume of blood to meet the body's needs), fainting with exertion (known as “exertional syncope”), or sudden death
- Abdominal swelling or distension
- Difficulty breathing (known as “dyspnea”)
- Exercise intolerance
- Heart murmur
- May be able to feel vibrations caused by abnormal blood flow (known as “thrills”) when placing hand against the chest wall (generally associated with more severe narrowing [stenosis])
- Irregular heartbeats (known as “arrhythmias”)
- Rapid heart rate (known as “tachycardia”) may occur if pet is in congestive heart failure (condition in which the heart cannot pump an adequate volume of blood to meet the body's needs)
- Other signs of congestive heart failure include fluid buildup in the abdomen (known as “ascites”), enlargement of the jugular veins in the neck, and rapid breathing (known as “tachypnea”)

CAUSES

- Congenital (present at birth) heart defect

Treatment

HEALTH CARE

- Most managed as outpatients
- Initial hospitalization of those pets with severe congestive heart failure; “congestive heart failure” is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Rarely, fluid buildup in the space between the lungs and chest wall (known as “pleural effusion”) may need draining; fluid buildup in the abdomen (ascites) usually is treated medically

ACTIVITY

- Exercise should be restricted in pets with fainting (syncope) or congestive heart failure (condition in which the heart cannot pump an adequate volume of blood to meet the body's needs)
- Excessive exertion should be avoided in pets with severe narrowing (stenosis) that do not have clinical signs

DIET

- Low-salt diets may benefit those pets with fluid buildup in the abdomen that does not respond to medical treatment (known as “refractory ascites”)

SURGERY

- Balloon dilation (procedure in which an instrument with an expandable balloon is inserted into the area of the narrowing [stenosis] and the balloon is expanded to open the narrowing) of the outflow tract, performed during heart catheterization
- Alternative surgical techniques include cutting through the narrowed pulmonary valve to relieve blockage to blood flow (procedure known as a “valvulotomy”) or patch-graft procedures; mortality rates tend to be higher than with balloon dilation

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive.

- If pet has signs of congestive heart failure, treat fluid buildup in the abdomen (ascites) with medications to remove excess fluids from the body (known as “diuretics”), such as furosemide or spironolactone (used as an additional diuretic in cases that do not respond to medical treatment); treat atrial fibrillation (rapid, irregular

heart rhythm involving the top two chambers of the heart [atria]) with digoxin

- Medications to enlarge or dilate blood vessels (known as “vasodilators,” such as hydralazine) may cause low blood pressure (known as “hypotension”) without relieving the narrowing (stenosis) and are best avoided

Follow-Up Care

PATIENT MONITORING

- Use serial follow-up echocardiograms (use of ultrasound to evaluate the heart and major blood vessels)

PREVENTIONS AND AVOIDANCE

- Do not breed affected pets

POSSIBLE COMPLICATIONS

- Right-sided congestive heart failure; congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Irregular heartbeats (arrhythmias)
- Exercise intolerance
- Fainting with exertion (exertional syncope)
- Sudden death

EXPECTED COURSE AND PROGNOSIS

- Mildly affected pets may remain without clinical signs and have normal life spans
- Severely affected pets have a guarded prognosis because they may develop congestive heart failure (condition in which the heart cannot pump an adequate volume of blood to meet the body's needs) or sudden death
- Clinical signs generally are more common in pets over 1 year of age

Key Points

- Mildly affected pets may lead normal lives
- Moderately to severely affected pets may benefit from treatment interventions (such as balloon catheter dilation or surgery); improved clinical signs and survival have been associated with successful balloon procedures
- Prognosis is guarded once signs of congestive heart failure develop; congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Do not breed affected pets

Notes

Enter notes here



Blackwell's Five-Minute Veterinary Consult: Canine and Feline, Fifth Edition, Larry P. Tilley and Francis W.K. Smith, Jr. © 2011 John Wiley & Sons, Inc.