

WOODVALE PARK

VETERINARY HOSPITAL



Unit 10, 923 Whitfords Avenue, Woodvale WA 6026

Phone: (08) 9409 6968

www.woodvaleparkvet.com.au

Aimeroy Pty Ltd ABN 53 165 893 701

Atrial Standstill

Basics

OVERVIEW

- The heart of the dog or cat is composed of four chambers; the top two chambers are the right and left atria and the bottom two chambers are the right and left ventricles
- “Atrial” refers to the atrium (singular) or atria (plural) of the heart
- “Atrial standstill” is an abnormal heart rhythm seen on a recording of an electrocardiogram (ECG, a recording of the electrical activity of the heart), characterized by absence of P waves—the P wave is the first upward deflection of the ECG tracing that looks like a “bump” in the tracing; the P wave is a measure of the electrical activity of the atria—absence of P waves indicates an absence of atrial activity; atrial standstill can be temporary (such as associated with too high levels of potassium in the blood [known as “hyperkalemia”] or induced by medications); persistent; or terminal (such as associated with severely high levels of potassium in the blood [severe hyperkalemia] or dying heart)
- Rare heart rhythm disturbance

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs
- Cats

Breed Predispositions

- Persistent atrial standstill—most common in English springer spaniels; other breeds occasionally affected

Mean Age and Range

- Most pets with persistent atrial standstill are young
- Pets with inadequate production of steroids by the adrenal glands (known as “hypoadrenocorticism” or “Addison’s disease”) leading to too high levels of potassium in the blood (hyperkalemia) are usually young to middle-aged

Predominant Sex

- Inadequate production of steroids by the adrenal glands (hypoadrenocorticism or Addison’s disease) more common in females (69%)

SIGNS/OBSERVED CHANGES IN THE PET

- Vary with underlying cause
- Sluggishness (lethargy) is common; fainting (known as “syncope”) may occur

- Pets with persistent atrial standstill may show signs of congestive heart failure (CHF); CHF signs include cough, difficulty breathing (dyspnea), bluish discoloration of the skin and moist tissues (mucous membranes) of the body caused by inadequate oxygen levels in the red blood cells (known as “cyanosis”); congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body’s needs
- Slow heart rate (known as “bradycardia”) is common
- Pets with persistent atrial standstill may have skeletal muscle wasting of the forearms and shoulders

CAUSES

- Too high levels of potassium in the blood (hyperkalemia)
- Heart disease involving the upper chambers of the heart (known as “atrial disease”), often associated with dilated or distended atria (such as cats with disease of the heart muscle [known as “cardiomyopathy”])
- Muscle disorder of the atria (known as “atrial myopathy”)—persistent atrial standstill

RISK FACTORS

Atrial Standstill Due to Too High Levels of Potassium in the Blood (Hyperkalemia)

- Inadequate production of steroids by the adrenal glands (hypoadrenocorticism or Addison’s disease) leading to increased levels of potassium in the blood
- Conditions leading to obstruction or rupture of the urinary tract
- Kidney failure characterized by production of only small amounts of urine (known as “oliguria”) or no urine (known as “anuria”)

Treatment

HEALTH CARE

Persistent Atrial Standstill

- Not life-threatening condition; pet can be treated as an outpatient

Atrial Standstill Due to Too High Levels of Potassium in the Blood (Hyperkalemia)

- Potentially life-threatening; often requires aggressive treatment
- Aggressive fluid therapy with 0.9% saline often necessary to correct low circulating blood volume (known as “hypovolemia”) and to lower serum potassium levels in pets with atrial standstill due to too high levels of potassium in the blood (hyperkalemia)

ACTIVITY

- Restrict activity in pets with persistent atrial standstill and signs of congestive heart failure or fainting (syncope); CHF signs include cough, difficulty breathing (dyspnea), bluish discoloration of the skin and moist tissues (mucous membranes) of the body caused by inadequate oxygen levels in the red blood cells (known as “cyanosis”); congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body’s needs

SURGERY

Persistent Atrial Standstill

- Surgically implant a permanent ventricular pacemaker to regulate heart rate and rhythm

Atrial Standstill Due to Too High Levels of Potassium in the Blood (Hyperkalemia)

- Too high levels of potassium in the blood (hyperkalemia) secondary to urinary tract obstruction or rupture may require surgery to treat the problems in the urinary tract

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive

Persistent Atrial Standstill

- Treat with diuretics to remove excess fluid from the body (such as furosemide or spironolactone) and angiotensin converting enzyme inhibitors (ACE inhibitors) to dilate blood vessels (such as enalapril or benazepril), if congestive heart failure develops

Atrial Standstill Due to Too High Levels of Potassium in the Blood (Hyperkalemia)

- Treat the underlying cause (such as kidney failure characterized by production of only small amounts of urine [oliguria], inadequate production of steroids by the adrenal glands [hypoadrenocorticism or Addison's disease])
- Aggressive fluid therapy with 0.9% saline and possibly sodium bicarbonate or insulin with dextrose
- Calcium gluconate—counters the heart effects of too high levels of potassium in the blood (hyperkalemia); can be used in life-threatening situations to reestablish a normal heart rhythm while instituting treatment to lower potassium concentration in the blood

Follow-Up Care

PATIENT MONITORING

- Monitor electrocardiogram (ECG, a recording of the electrical activity of the heart) during treatment of too high levels of potassium in the blood (hyperkalemia) and periodically in pets with a permanent ventricular pacemaker
- Monitor electrolytes in pets with atrial standstill due to too high levels of potassium in the blood (hyperkalemia)
- Monitor pets with persistent atrial standstill for signs of congestive heart failure; CHF signs include cough, difficulty breathing (dyspnea), bluish discoloration of the skin and moist tissues (mucous membranes) of the body caused by inadequate oxygen levels in the red-blood cells (known as “cyanosis”); congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs

POSSIBLE COMPLICATIONS

- Congestive heart failure in pets with persistent atrial standstill

EXPECTED COURSE AND PROGNOSIS

Persistent Atrial Standstill

- Clinical signs generally improve after pacemaker implantation
- Signs of congestive heart failure may develop, and weakness and sluggishness (lethargy) may persist even after heart rate and rhythm are corrected with the pacemaker; CHF signs include cough, difficulty breathing (dyspnea), bluish discoloration of the skin and moist tissues (mucous membranes) of the body caused by inadequate oxygen levels in the red blood cells (known as “cyanosis”); congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs

Atrial Standstill Due to Too High Levels of Potassium in the Blood (Hyperkalemia)

- Long-term prognosis is excellent if underlying cause can be corrected and the increased levels of potassium in the blood (hyperkalemia) are reversed

Key Points

PERSISTENT ATRIAL STANDSTILL

- Clinical signs generally improve after pacemaker implantation
- Signs of congestive heart failure may develop, and weakness and sluggishness (lethargy) may persist even after heart rate and rhythm are corrected with the pacemaker; CHF signs include cough, difficulty breathing (dyspnea), bluish discoloration of the skin and moist tissues (mucous membranes) of the body caused by inadequate oxygen levels in the red blood cells (known as “cyanosis”); congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs

ATRIAL STANDSTILL DUE TO TOO HIGH LEVELS OF POTASSIUM IN THE BLOOD (HYPERKALEMIA)

- Potentially life-threatening; often requires aggressive treatment

Notes

Enter notes here



Blackwell's Five-Minute Veterinary Consult: Canine and Feline, Fifth Edition, Larry P. Tilley and Francis W.K. Smith, Jr. © 2011 John Wiley & Sons, Inc.