

WOODVALE PARK

VETERINARY HOSPITAL



Unit 10, 923 Whitfords Avenue, Woodvale WA 6026

Phone: (08) 9409 6968

www.woodvaleparkvet.com.au

Aimeroy Pty Ltd ABN 53 165 893 701

Pneumothorax

(Air in the Pleural Space, the Space between the Chest Wall and the Lung)

Basics

OVERVIEW

- Air accumulation in the pleural space (the space between the chest wall and the lungs); it is categorized as “traumatic” or “spontaneous” and “closed” or “open”
- “Traumatic” pneumothorax—air accumulates in the pleural space, following some type of trauma (such as being hit by a car)
- “Spontaneous” pneumothorax—air accumulates in the pleural space in the absence of trauma ; it is a “closed” pneumothorax
- “Closed” pneumothorax—no defects (such as a puncture) in the chest wall; air leaks into the pleural space from the lungs or large airways
- “Open” pneumothorax—defect (such as a puncture) in the chest wall, resulting in communication of the pleural space (the space between the chest wall and the lungs) with the atmosphere
- “Tension” pneumothorax—air is transferred into the pleural space (the space between the chest wall and the lungs) during inspiration and it becomes trapped, condition creates a one-way transfer of air into the pleural space from room or outdoor air

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs
- Cats

Breed Predispositions

- Spontaneous pneumothorax—more common in large, deep-chested dogs; Siberian huskies may be more susceptible than other dog breeds

SIGNS/OBSERVED CHANGES IN THE PET

- Traumatic pneumothorax—evidence of recent trauma
- Spontaneous pneumothorax—may or may not have previous signs of lung disease; usually sudden (acute), but can have a slowly progressive onset
- Rapid breathing (known as “tachypnea”)
- Difficulty breathing (known as “dyspnea”)

- Respiratory distress
- Standing with the elbows away from the body in an attempt to increase lung capacity (known as “orthopnea”)
- Shallow, rapid abdominal breathing common
- Rapid heart rate (known as “tachycardia”)
- Reduced lung sounds near the back—can be difficult to appreciate in pets with severe difficulty breathing

Traumatic Pneumothorax

- Other signs of trauma, including pale gums and moist tissues of the body (known as “mucous membranes”); bluish discoloration of the skin and mucous membranes caused by inadequate oxygen levels in the red blood cells (condition known as “cyanosis”) in severely affected pets; shock, where circulation or flow of blood is unable to sustain the body
- May or may not have evidence of chest trauma
- Open pneumothorax—obvious chest-wall trauma present
- Air under the skin (known as “subcutaneous emphysema”) in some cases with air in the mediastinum (the mediastinum is the center portion of the chest that contains the heart and other organs [except for the lungs]; condition is known as “pneumomediastinum”) and/or trauma to the windpipe or trachea

CAUSES

- Traumatic pneumothorax—blunt trauma; penetrating chest injuries; penetrating injuries to the neck; following medical procedure to tap the chest (known as “thoracocentesis”); following surgical incision into the chest (known as “thoracotomy”); perforation of the esophagus (the tube running from the throat to the stomach); injury to the windpipe (trachea) following endotracheal intubation during anesthesia/surgery
- Spontaneous pneumothorax—condition characterized by enlarged or dilated airspaces (known as “bullous emphysema”) is most common cause in dogs; migrating foreign body in the lung; lung cancer, lung abscess; feline asthma; pneumonia; fungal lung nodules (known as “mycotic pulmonary granulomas”); lung disease caused by parasites (such as *Paragonimus*); blister or bubble-like structures in the lungs (known as “pulmonary bullae”); large air-filled or fluid-filled sacs in the lungs (known as “pulmonary blebs”)

RISK FACTORS

- Trauma
- Tapping the chest, to withdraw fluid or air (thoracocentesis)
- Surgical incision into the chest (thoracotomy)
- Overinflating the cuff of endotracheal tube, during anesthesia/surgery
- Lung disease
- Migrating grass awns

Treatment

HEALTH CARE

- Inpatient care, until air accumulation has stopped or has stabilized
- Pets with respiratory distress should have the chest tapped (thoracocentesis) to remove a maximal amount of air
- Provide oxygen therapy, until the pet is stabilized
- Pain relief (analgesia) with an opioid-type drug, if significant injuries following trauma
- Tapping the chest (thoracocentesis) can be performed with an intravenous catheter attached to an extension set and stopcock or via a butterfly needle; no need to perform thoracocentesis on pet that is not having breathing difficulties; if large open chest wound—cover as cleanly as possible with airtight bandage (use of sterile lubricant/ointment around periphery of wound); must be accompanied by chest-tube placement; will require surgical closure of chest wound, once pet is stable
- Intravenous fluids required in most cases of trauma
- Chest-tube maintenance—ensure all connections are air-tight (cable ties are excellent for securing connections); ensure that tube is attached to pet at two points to reduce chance of inadvertent tube removal; clean tube site and change dressing once daily—do not allow pet to chew at chest tube

ACTIVITY

- Strict rest for at least a week following resolution of pneumothorax, in an effort to minimize the chance of

recurrence

SURGERY

- Establishing an opening into the chest with a tube (known as “tube thoracostomy”)—use if unable to stabilize with tapping the chest (thoracocentesis) or repeated thoracocentesis procedures are required for continued pneumothorax (air in the space between the chest and the lungs); chest-tube placement (under local or general anesthesia)—chest tube is passed into pleural space, and a purse-string suture is placed in the skin and the tube is secured with “finger-trap” suture pattern; chest x-rays (radiographs) should be performed after chest-tube placement
- If air is accumulating rapidly in the space between the chest wall and lungs (pneumothorax)—use continuous chest-tube suction; if pneumothorax is not severe or is resolving—use intermittent chest-tube aspiration
- In emergency situation of life-threatening tension pneumothorax (in which air is transferred into the pleural space [the space between the chest wall and the lungs] during inspiration and becomes trapped, creating one-way transfer of air into the pleural space)—consider emergency surgical incision into the chest (known as “thoracotomy”) to convert problem to an “open pneumothorax”; pet then can be intubated with an endotracheal tube and breathing can be controlled by an assistant or by a ventilator, until the pet is stabilized
- Open traumatic pneumothorax—surgery as soon as the pet is stable
- Closed traumatic pneumothorax—rarely requires surgical intervention
- Spontaneous pneumothorax—early surgical intervention recommended in dogs; exploratory surgical incision into the chest (thoracotomy) often performed to allow visualization of the lungs, to determine location of the air leakage
- Evaluation of the space between the chest wall and lungs with an endoscope (a special lighted instrument; procedure known as “thoracoscopy”)—may allow visualization of local lesion; allows instillation of substances to cause formation of scar tissue across the pleural space (the space between the chest wall and the lungs) to eliminate the space where air has been collecting (procedure is known as “pleurodesis”)
- Removal of part or all of a lung may be necessary for localized lesions; traumatic lacerations may be sutured
- Pleurodesis (to cause formation of scar tissue across the pleural space [the space between the chest wall and the lungs] to eliminate the space where air has been collecting) with mechanical abrasion of the pleura (lining of the chest) or instillation of a substance intended to cause inflammation in the pleural space

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive

- Medications to control pain (known as “analgesics”) should be used carefully; some analgesics can decrease breathing effort

Follow-Up Care

PATIENT MONITORING

- Breathing rate—increased rate suggests recurrence of pneumothorax (free air in the space between the chest wall and lungs)
- Repeated chest x-rays (radiographs) to determine amount of air accumulating
- Pulse oximetry (a means of measuring oxygen levels in blood), if breathing room air can help determine oxygenation status; arterial blood gases (measurements of oxygen and carbon dioxide levels in arterial blood) give the best evaluation of oxygenation status, if lung disease is present
- Rate of air removed from chest through the chest tube

PREVENTIONS AND AVOIDANCE

- Keep pets confined—less likely to be injured

POSSIBLE COMPLICATIONS

- Death from low levels of oxygen in the blood (known as “hypoxemia”) and impairment of the circulation or heart and blood vessels (known as “cardiovascular compromise”)

- Incorrect placement of chest tube or trauma associated with tapping the chest (thoracocentesis) —injury to other organs of the body, such as lung-lobe laceration, heart puncture, diaphragmatic laceration, liver trauma
- Infection of the lining of the chest (known as “pleural infection”) from tapping the chest (thoracocentesis) or presence of the chest drain

EXPECTED COURSE AND PROGNOSIS

- Traumatic pneumothorax—if chest trauma is not severe, the prognosis is good with tapping the chest to remove the air (thoracocentesis) and/or chest-drain placement
- Traumatic pneumothorax—with severe chest trauma, the pet can deteriorate despite all efforts to stabilize it, usually because of severe bruising of the lungs and/or bleeding into the lungs
- Spontaneous pneumothorax—prognosis depends on underlying cause; if a single, localized lesion can be removed surgically, prognosis is good; if unable to locate lesion or generalized lung disease is present—prognosis is poor

Key Points

- Traumatic pneumothorax—possibility of a chest tube and need for hospitalization; some pets may require surgery
- Spontaneous pneumothorax—recommend early surgical intervention in most cases in dogs; possibility of underlying lung disease that may make resolution challenging and recurrence possible—even with surgical incision into the chest (thoracotomy), the source of the leaking air (pneumothorax) may not be found; recurrence of spontaneous pneumothorax is possible

Notes

Enter notes here

