



Restrictive Cardiomyopathy in Cats

(a Type of Heart-Muscle Disease)

Basics

OVERVIEW

- The heart of the cat is composed of four chambers; the top two chambers are the left and right atria and the bottom two chambers are the left and right ventricles; heart valves are located between the left atrium and the left ventricle (mitral valve); between the right atrium and the right ventricle (tricuspid valve); from the left ventricle to the aorta (the main artery of the body; valve is the aortic valve); and from the right ventricle to the main pulmonary (lung) artery (pulmonary valve)
- “Cardiomyopathy” is the medical term for disease of the heart muscle; “restrictive cardiomyopathy” is a disease in which the muscle is “stiff” and does not expand, such that blood cannot fill the ventricles normally
- “Restrictive cardiomyopathy” in cats is characterized by abnormal filling of the chambers of the heart (known as “diastolic dysfunction”), severe atrial enlargement, normal left ventricular wall thickness and variable abnormal pumping of the heart (known as “systolic dysfunction”)
- Scar tissue of the heart muscle layer may be present; in addition, other changes or damage in the muscle may be associated with other heart-muscle disorders, including inflammatory or immune-mediated diseases

SIGNALMENT/DESCRIPTION OF PET

Species

- Cats

Mean Age and Range

- Middle-aged to older cats

SIGNS/OBSERVED CHANGES IN THE PET

If Cat Does Not Have Congestive Heart Failure

- Congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Some cats have no clinical signs
- Sluggishness (lethargy)
- Poor appetite and weight loss

- Fainting (known as “syncope”)—rare; usually indicates serious irregular heartbeats (known as “arrhythmias”)
- Weakness or paralysis (signs of blockage of blood flow secondary to the presence of a blood clot in the artery [condition known as “arterial thromboembolism”])
- Depression
- Extreme weight loss with muscle wasting (known as “cachexia”)
- Rapid heart rate (known as “tachycardia”)
- Irregular heartbeats (arrhythmias)
- Sequence of three heart sounds (known as a “gallop rhythm”), when listening to the heart with a stethoscope; heart beat sounds like a galloping horse instead of normal “lub-dub”
- May have a heart murmur

If Cat Has Congestive Heart Failure

- Congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Cat has signs as previously described, plus the following:
 - ♦ Difficulty breathing (known as “dyspnea”)
 - ♦ Rapid breathing (known as “tachypnea”)
 - ♦ Panting
 - ♦ Open-mouth breathing
 - ♦ Bluish discoloration of the skin and moist tissues (known as “mucous membranes”) of the body caused by inadequate oxygen levels in the red blood cells (known as “cyanosis”)
 - ♦ Abdominal swelling or distention
 - ♦ Enlarged liver (known as “hepatomegaly”) or fluid buildup in the abdomen (known as “ascites”), with enlargement or distension of the jugular veins (located on either side of the neck)
 - ♦ Short, rough snapping sounds (known as “crackles”) heard when listening to the chest with a stethoscope
 - ♦ Muffled heart or lung sounds heard when listening to the chest with a stethoscope, if the cat has fluid buildup in the space between the chest wall and lungs (known as “pleural effusion”)
 - ♦ Weakness or paralysis with loss of femoral pulses; one or more extremities may be cold and painful (signs of blockage of blood flow secondary to the presence of a blood clot in the artery [condition is “arterial thromboembolism”])

CAUSES

- True cause(s) unknown (so-called “idiopathic disease”); often no “predisposing” disease can be documented
- Suspected initiating causes include inflammation of the heart muscle (known as “myocarditis”); inflammation of the inner muscle layer of the heart (known as “endomyocarditis”); infiltration of eosinophils (a type of white-blood cell) into the heart muscle (known as “eosinophilic myocardial infiltration”); disease characterized by inappropriate enlargement or thickening of the heart muscle of the left ventricle (known as “hypertrophic cardiomyopathy”) with sudden lack of blood supply to the heart muscle that leads to death of tissues (known as “myocardial infarction”); widespread (diffuse) “small blood vessel disease;” and other causes of abnormal blood flow and resulting lack of oxygen to the heart muscle

Treatment

HEALTH CARE

- Pets with sudden (acute), severe congestive heart failure are hospitalized for emergency care; “congestive heart failure” is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Pets that do not have clinical signs or have mild signs can be treated with outpatient medical management
- Pets with severe difficulty breathing (severe dyspnea) should receive oxygen via oxygen cage, nasal cannula, or mask (beware of stress to the pet)
- Life-threatening fluid buildup in the space between the chest wall and lungs (pleural effusion) is reduced via tapping and draining the chest (known as “thoracocentesis”); perform thoracocentesis as necessary to relieve difficulty breathing (dyspnea)
- Treat associated conditions (such as dehydration or low body temperature [hypothermia])

- Low sodium fluids administered cautiously if dehydration occurs (beware of worsening congestive heart failure)
- Heating pad may be necessary for pets with low body temperature (hypothermia)

ACTIVITY

- Maintain a low-stress environment to decrease pet anxiety (such as cage rest, minimize handling)
- Most cats in congestive heart failure will limit their own exercise; congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs

DIET

Medications

- Low-salt diet may decrease fluid retention, but strict adherence to dietary changes should be avoided in sudden (acute) congestive heart failure in order to maintain food intake
- Hand-feed, as necessary

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive

SUDDEN (ACUTE) CONGESTIVE HEART FAILURE

- Congestive heart failure is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Medication to remove excess fluid from the body (diuretic) administered by injection—furosemide
- Dobutamine (drug to increase contraction of the heart muscle) to increase function of the heart
- Nitroglycerin ointment, applied to the skin
- Oxygen delivered by cage, mask, or nasal tube
- Severe irregular heartbeats originating above the ventricles (supraventricular arrhythmias) may be treated with diltiazem
- Rapid ventricular heart rate (ventricular tachycardia) may resolve with resolution of congestive heart failure; treatment for sudden occurrence of ventricular tachycardia may include lidocaine
- Beta-blockers (such as propranolol or atenolol) may be used to treat irregular heartbeats that originate above or in the ventricles (supraventricular or ventricular arrhythmias), but not until congestive heart failure is treated
- Pimobendan may be helpful to improve heart function; it is not approved for use in cats

LONG-TERM (CHRONIC) THERAPY

- Medication to remove excess fluid from the body (diuretic)—furosemide, gradually decreased to lowest effective dose
- Long-term (chronic) therapy with diltiazem decreases heart rate and improves irregular heartbeats that originate above the ventricles (supraventricular arrhythmias)
- Beta-blockers may be used to slow heart rate and treat irregular heartbeats that originate above or in the ventricles (supraventricular or ventricular arrhythmias)
- Angiotensin converting enzyme (ACE) inhibitors may reduce fluid retention and decrease need for medications to remove excess fluid from the body (diuretics); examples of ACE inhibitors are enalapril and benazepril
- Pimobendan may be helpful to improve heart function; it is not approved for use in cats
- Digoxin (a heart medication) may be used if heart-muscle contraction is impaired or atrial fibrillation (rapid, irregular heart rhythm involving the top two chambers of the heart [atria]) is present
- Aspirin may be administered to prevent blood clots (known as “thromboembolism”), but effectiveness is questionable; administer aspirin only under the direction of your cat's veterinarian
- Warfarin may be administered to prevent blood clots (thromboembolism), but is not recommended unless close monitoring and repeated measurement of prothrombin time (a blood test to evaluate clotting) are feasible
- Clopidogrel may be used to decrease clumping of platelets; effectiveness is not proven

Follow-Up Care

PATIENT MONITORING

- Frequent serial physical examinations (minimal stress to the pet) to assess response to treatment and resolution of fluid buildup in the lungs (pulmonary edema) and fluid buildup in body cavities (known as “effusions”)
- Monitor breathing rate to assess response to removing fluid from the space between the chest wall and lungs (pleural effusion) via tapping and draining the chest (thoracocentesis) or administration of medications to decrease fluid buildup in the body (diuretics); breathing rate should decrease with successful treatment
- Frequent assessment of hydration and kidney function is important in first few days of therapy to avoid removal of too much fluid from the body (known as “over diuresis”) and development of excessive levels of urea and other nitrogenous waste products in the blood (known as “uremia” or “azotemia”)
- Repeated tapping and draining the chest (thoracocentesis) may be necessary to maintain the amount of fluid buildup in the space between the chest wall and lungs (pleural effusion) at a comfortable level
- “Hands-off” hourly assessment of breathing rate in first 12–24 hours can be used to monitor effectiveness of congestive heart failure therapy
- Chest x-rays (radiographs) may be repeated in 12–24 hours
- Blood work (especially creatinine and potassium) should be monitored closely during the first 3–5 days of therapy to detect dehydration, kidney failure, and low levels of potassium in the blood (known as “hypokalemia”)—caused by medications to remove excess fluid from the body (diuretics); or high levels of potassium in the blood (known as “hyperkalemia”), if angiotensin converting enzyme inhibitors are administered
- Repeat physical examination and blood work (especially electrolyte analysis) after approximately 10–14 days of treatment
- Electrocardiograms (ECGs, recordings of the electrical activity of the heart) and x-rays (radiographs) may be repeated, as your pet's veterinarian feels necessary
- Stable pets are reevaluated every 2–4 months, or more frequently if problems occur

PREVENTIONS AND AVOIDANCE

- No known prevention of restrictive cardiomyopathy in cats
- Medications to prevent clumping of platelets (such as aspirin or clopidogrel) or to prevent blood clotting (such as warfarin) have been recommended to prevent complications from blood clots; effectiveness of such treatment is not proven

POSSIBLE COMPLICATIONS

- Congestive heart failure
- Loss of function of the legs or death of tissues due to lack of circulation (complication from blood clots secondary to restrictive cardiomyopathy)
- Death

EXPECTED COURSE AND PROGNOSIS

- Highly variable, based on presentation of disease and clinical signs
- Most cats with restrictive cardiomyopathy and congestive heart failure live 3–12 months; some live 2 years

Key Points

- “Restrictive cardiomyopathy” is a disease in which the heart muscle is “stiff” and does not expand, such that blood cannot fill the ventricles normally
- Pets with sudden (acute), severe congestive heart failure are hospitalized for emergency care; “congestive heart failure” is a condition in which the heart cannot pump an adequate volume of blood to meet the body's needs
- Pets that do not have clinical signs or have mild signs can be treated with outpatient medical management
- Monitor your pet's breathing rate at home and keep a written tracker of breathing rates; contact your pet's veterinarian if the breathing rate increases suddenly or progressively over several days
- Monitor your pet's food and water intake at home; the goal is that the pet maintains a stable food and water intake
- Most cats with restrictive cardiomyopathy and congestive heart failure live 3–12 months; some live 2 years

Notes

Enter notes here



Blackwell's Five-Minute Veterinary Consult: Canine and Feline, Fifth Edition, Larry P. Tilley and Francis W.K. Smith, Jr. © 2011 John Wiley & Sons, Inc.