

WOODVALE PARK

VETERINARY HOSPITAL



Unit 10, 923 Whitfords Avenue, Woodvale WA 6026

Phone: (08) 9409 6968

www.woodvaleparkvet.com.au

Aimeroy Pty Ltd ABN 53 165 893 701

Vaginal Discharge in Dogs

Basics

OVERVIEW

- “Vaginal” refers to the vagina; the “vagina” is the tubular passageway leading from the opening of the vulva to the cervix of the uterus; “vulvar” refers to the vulva; the “vulva” is the external genitalia of females
- “Vaginal discharge” is any substance (such as blood, mucus, pus) coming from the vagina, through the vulvar opening
- “Bitch” is a female dog

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs

Mean Age and Range

- Bitches (less than 6–12 months of age) prior to going through puberty (known as “prepubertal bitches”)—anatomic abnormalities and prepubertal inflammation of the vagina (known as “prepubertal vaginitis” or “puppy vaginitis”) more common
- Bitches in “heat” or “estrus” or following delivery of puppies (whelping)—normal vaginal discharges unless they are discolored or are malodorous
- Bitches that recently have completed their “heat” or “estrous cycle” or are pregnant or following delivery of puppies (whelping)—vaginal discharge may be more serious

Predominant Sex

- Females

SIGNS/OBSERVED CHANGES IN THE PET

- Discharge from the vulva (the external genitalia); discharge may be blood; blood, mucus, and tissue debris (known as “lochia”) following delivery of puppies; pus; urine; or feces
- Licking the vulva (external female genitalia)
- Spotting
- Scooting
- Attracting male dogs
- Delivering puppies (whelping or parturition)—with post-partum discharge
- History of “heat” or “estrus” during the preceding 2 months—vaginal discharge may be related to inflammation with accumulation of pus in the uterus (known as “pyometra”)
- Vomiting and lack of appetite (known as “anorexia”)—may be seen with inflammation of the lining of the uterus (known as “metritis”) or inflammation with accumulation of pus in the uterus (known as “pyometra”)

CAUSES

Discharge Containing Serum and Blood (Known as “Serosanguineous Discharge”)

- Normal during early heat cycle (known as “proestrus”) and sometimes into “heat” or “estrus”
- Urinary tract infection
- Foreign body
- Tumor or cancer of the vagina—such as transmissible venereal tumor; leiomyoma
- Vaginal trauma
- Fetal death
- Localized accumulation of blood in the vagina (known as a “vaginal hematoma”)
- Tumor or cancer of the ovaries
- Blood-clotting disorders (known as “coagulopathies”)

Discharge Containing Blood, Mucus, and Tissue Debris (Lochia) and Other Discharges Following Birth (Post-partum)

- Normal post-partum discharge—for 6–8 weeks
- Subinvolution of placental sites (condition in which the placental attachment sites in the lining of the uterus do not return to normal in 6 to 8 weeks after delivery)—discharge lasting longer than 8 weeks
- Retained placentas or afterbirth (condition in which one or more placentas remains within the uterus following the delivery of the puppy or puppies)
- Inflammation of the lining of the uterus (metritis)

Discharge Containing Pus

- Normal in early diestrus (“diestrus” is the phase of the “heat” cycle immediately after “standing heat”; slight amount of discharge)
- Inflammation of the vagina in a bitch prior to puberty (prepubertal vaginitis)
- Primary inflammation of the vagina (vaginitis)
- Secondary inflammation of the vagina (vaginitis)—from anatomic abnormality, foreign body, urinary tract infection, enlargement of the clitoris (known as “clitoral hypertrophy”), vaginal tumor, and fetal death
- Inflammation with accumulation of pus in the uterus (pyometra)
- Embryonic and fetal death
- Inflammation of the lining of the uterus following birth of the puppies (post-partum metritis)
- Inflammation of the skin around the vulva (known as “perivulvar dermatitis”)
- Zinc toxicity—reported

Other

- Urine or feces—with congenital (present at birth) anatomic abnormalities
- Acquired (condition that develops sometime later in life/after birth) inflammation of the skin around the vulva (perivulvar dermatitis) also can be mistaken for vaginal discharge
- Urine from ectopic ureters or lack of control of urination (incontinence) from low levels of estrogen (known as “hypoestrogenism”); the ureters are tubes from the kidneys to the bladder; during development, they may not attach to the bladder properly or may attach to reproductive organs instead; when this occurs, they are called “ectopic ureters” and one or both can terminate in the lower urethra, uterus, or vagina
- Normal mucous discharge during pregnancy

RISK FACTORS

- Medications or products containing male hormones (androgens)—may cause enlargement of the clitoris (clitoral hypertrophy)
- Prophylactic antibiotics—may alter the normal vaginal bacteria and allow overgrowth of disease-causing species and increase likelihood of secondary infection
- Estrogen medications given during certain phases of the “heat” or “estrous” cycle—predispose the pet to inflammation with accumulation of pus in the uterus (pyometra)
- Obesity—excess skin folds around the vulva (female external genitalia)

Treatment

HEALTH CARE

- Based on cause

- No treatment needed for normal causes of vaginal discharge
- Outpatient, unless inflammation of the lining of the uterus (metritis) or inflammation with accumulation of pus in the uterus (pyometra) is noted and spaying (ovariohysterectomy) may be indicated
- Supportive fluids—for inflammation of the lining of the uterus (metritis) or inflammation with accumulation of pus in the uterus (pyometra), if the pet is ill
- Medical treatment for inflammation with accumulation of pus in the uterus (pyometra)—performed in a hospital and with great care
- Remove or treat any inciting cause—foreign body; tumor; anatomic abnormality; urinary tract infection; medications or products containing male hormones (androgens) or estrogens
- Inflammation of the vagina in a bitch prior to puberty (prepubertal vaginitis)—usually resolves spontaneously after the first “heat” or “estrus”
- Subinvolution of placental sites (condition in which the placental attachment sites in the lining of the uterus do not return to normal in 6–8 weeks)—rarely requires treatment

SURGERY

- Depends on cause of vaginal discharge
- Bitches with inflammation of the lining of the uterus (metritis) or inflammation with accumulation of pus in the uterus (pyometra) may require surgery; a “spay” or “ovariohysterectomy” (in which the ovaries and uterus are removed surgically) may be indicated
- Bitches with congenital (present at birth) anatomic abnormalities may need surgery
- Bitches with cancer—surgical removal of the ovary (known as “ovariectomy”) or the ovaries and uterus (ovariohysterectomy) is the treatment of choice
- Bitches with transmissible venereal tumor (TVT)—surgical removal or radiation therapy; “transmissible venereal tumor” is a tumor that is spread for dog to dog through physical contact, usually during breeding

Medications

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive

- Prostaglandin F_{2α} (PGF_{2α})—cause smooth muscle contractions and subsequent emptying of the uterus; also causes breakdown of the “corpora luteum” or “yellow body” that develops at the site of ovulation in the ovary and produces the female hormone, progesterone, which supports and maintains the pregnancy; breakdown or lysis of the corpora luteum decreases levels of progesterone and disrupts support of pregnancy
- Cloprostenol (Estrumate)—also used to breakdown the corpora luteum and decrease levels of progesterone
- Antibiotics—based on bacterial culture and sensitivity testing; culture should be obtained from deep in the vagina
- Bromocriptine or cabergoline—dopamine agonists: medications that mimic dopamine (a nervous system “messenger”) that leads to a decrease in the hormone, prolactin, and to lysis of the “corpora luteum” or “yellow body” via suppression of prolactin
- Aglepristone—may be used alone or with prostaglandin therapy for inflammation with accumulation of pus in the uterus (pyometra); currently not available in the United States

Follow-Up Care

PATIENT MONITORING

- Recheck serum progesterone levels until below 2.0 ng/ml; “progesterone” is the female hormone produced by the “corpus luteum” or “yellow body” in the ovary; it supports and maintains the pregnancy
- Ultrasonography or x-rays—after the progesterone drops below 2.0 ng/ml to monitor clearance of uterine fluid in pets having inflammation with accumulation of pus in the uterus (pyometra)
- Recheck bloodwork (complete blood count [CBC] and serum biochemistry profile) to monitor for generalized (systemic) health
- Recheck packed cell volume (PCV, a means of measuring the percentage volume of red blood cells as compared to the fluid volume of blood) for pets with subinvolution of placental sites to determine blood loss; subinvolution

of placental sites is the condition in which the placental attachment sites in the lining of the uterus do not return to normal in 6 to 8 weeks after delivery

PREVENTIONS AND AVOIDANCE

- Depends on cause of vaginal discharge
- Juvenile or puppy vaginitis—delay elective surgical removal of the ovaries and uterus (ovariohysterectomy) until after the first heat or estrous cycle; may avoid long-term (chronic) inflammation of the vagina (vagnitis)
- Avoid use of medications containing female hormones (estrogens or progestins) and male hormones (androgens) in intact bitches; an “intact female” has her reproductive organs and is capable of reproducing
- Spaying (ovariohysterectomy) will prevent inflammation of the lining of the uterus (metritis) or inflammation with accumulation of pus in the uterus (pyometra)

POSSIBLE COMPLICATIONS

- Bacterial toxins that accumulate in the blood (known as “endotoxemia”) and generalized disease caused by the spread of bacteria in the blood (known as “septicemia” or “blood poisoning”)—with inflammation of the lining of the uterus (metritis) or inflammation with accumulation of pus in the uterus (pyometra)

EXPECTED COURSE AND PROGNOSIS

- Depends on cause of vaginal discharge
- Grave prognosis for future fertility in bitches diagnosed with inflammation with accumulation of pus in the uterus (pyometra) in conjunction with cystic endometrial hyperplasia (CEH), a condition in which the lining of the uterus thickens abnormally and contains fluid-filled sacs or cysts; better prognosis if overt signs of CEH are not present and bitch is bred on next heat or estrous cycle
- Cancer—poor prognosis for future fertility
- Transmissible venereal tumor (TVT)—sexually transmitted disease; breeding should be avoided
- *Brucella canis* infection—sexually transmitted bacterial infection; grave prognosis for resolution of disease and normal fertility; affected pet should not be used for breeding; zoonotic potential (zoonoses are diseases that can be spread from animals to people)

Key Points

- Vaginal discharge is any substance (such as blood, mucus, pus) coming from the vagina, through the vulvar opening

Notes

Enter notes here



Blackwell's Five-Minute Veterinary Consult: Canine and Feline, Fifth Edition, Larry P. Tilley and Francis W.K. Smith, Jr. © 2011 John Wiley & Sons, Inc.